

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) Publication number:

0 473 159 B1

(12)

EUROPEAN PATENT SPECIFICATION

- (45) Date of publication of patent specification: **08.03.95** (51) Int. Cl.⁶: **A61K 9/06, A61K 31/07, A61K 31/23**
- (21) Application number: **91114550.6**
- (22) Date of filing: **29.08.91**

(54) **Stable aqueous preparation.**

- (30) Priority: **30.08.90 JP 229888/90**
- (43) Date of publication of application:
04.03.92 Bulletin 92/10
- (45) Publication of the grant of the patent:
08.03.95 Bulletin 95/10
- (84) Designated Contracting States:
AT BE CH DE DK ES FR GB GR IT LI LU NL SE
- (56) References cited:
EP-A- 0 258 865
US-A- 4 199 567
US-A- 4 966 773

PATENT ABSTRACTS OF JAPAN, vol. 7, no. 147 (C-173)[1292], 28th June 1983; & JP-A-58 59 257 (RAION K.K.) 08-04-1983

WPIL, FILE SUPPLIER, accession no. 84-058476 [10], Derwent Publications Ltd, London, GB; & JP-A-59 015 018 (JAPAN STORAGE BATTERY) 26-10-1984

(73) Proprietor: **Senju Pharmaceutical Co., Ltd.**
5-8, Hiranomachi 2-chome,
Chuo-Ku
Osaka-shi,
Osaka 541 (JP)

(72) Inventor: **Ushio, Kazumichi**
6-13 Kamiochi 1-chome
Nishinomiya,
Hyogo-ken (JP)
Inventor: **Yoshida, Kenichi**
103 Corpo Miyanoshita 2,
Kitahonmachi 2-chome
Itami,
Hyogo-ken (JP)

(74) Representative: **Kinzebach, Werner, Dr. et al**
Patentanwälte
Reitsötter, Kinzebach und Partner
Postfach 86 06 49
D-81633 München (DE)

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid (Art. 99(1) European patent convention).

EP 0 473 159 B1

D scription**BACKGROUND OF THE INVENTION**

5 The present invention relates to an aqueous preparation containing vitamin A or its derivative which may be used for the treatment of dry eye syndrome represented by lacrimal hyposecretion, squamous metaplasia and the like. More specifically the present invention relates to an aqueous preparation containing vitamin A or its derivative and having a stability which may allow the use of the preparation as an ophthalmic pharmaceutical preparation.

10 The cornea is an avascular tissue forming the front surface of the eyeball, and it has transparency and a constant curvature which are necessary to correctly introduce light from the environment into the eyeball. To maintain transparency of the cornea, it is the fundamental requirement that each of the 5 layers constituting the cornea, i.e. the corneal epithelium, the Bowman's membrane, the corneal stroma, the Descemet's membrane and the corneal endothelium, is regularly arranged and performs its normal
15 physiological functions. Since the cornea is an avascular tissue, its supplies of nutrients required for the maintenance of its functions are dependent on the perilimbal capillary loops, the aqueous humor and the tear fluid, and the corneal epithelium, in particular, is exclusively dependent on the tear fluid for the nutrient supplies. In addition, the corneal epithelium is protected by the tear film from the air and the mechanical stimulation which may be caused by friction with the palpebral conjunctiva.

20 Since the corneal epithelium is constantly exposed to the air except when the eyelids are closed, the water in the tear fluid which covers the cornea is constantly lost by evaporation, and the loss is supplemented by a continuous production of the tear fluid. The production of the tear fluid is made cooperated by the lacrimal gland, the goblet cells and the meibomian glands. Their functions may be lowered by a variety of extraocular disorders and systemic disorders such as autoimmune diseases as well
25 as aging. When the function to produce the tear fluid is lowered, the corneal epithelium tends to be dried, the physiological state of the corneal epithelium is disturbed by this, and the occurrence of inflammation is promoted, which may lead to corneal injury, corneal opacification or vascular infiltration into the cornea. Moreover, it is known that since the cornea is an extremely sensitive site, only a very slight tendency to dryness may cause uncomfortable sensation such as photophobia, pain, feeling of foreign body and the
30 like, thus hindering daily life, and may cause severe symptoms in some cases.

Vitamin A or its derivatives are known to be effective on such pathological states [e.g. J.L. Ubels et al., Current Eye Research, 10 (4), 1049 (1985)], and based on these reports, topical preparations which contain an effective amount of vitamin A or its derivative having vitamin A activities (in the description hereinafter, the simple wording "vitamin A" refers to "vitamin A and its derivatives having vitamin A activities", unless
35 derivatives of vitamin A are clearly excluded according to the context) have come to be applied. Among others, preparations in the form of aqueous eye drops have been needed so as to allow appropriate administration by patients themselves, since it is necessary to repeatedly administer vitamin A for certain period of time in order to treat patients with a variety of uncomfortable symptoms and severe damages caused by the tendency to dryness of the cornea and conjunctiva.

40 However, vitamin A and its derivatives are extremely unstable compounds. It is known that, inter alia, in aqueous preparations, their loss is likely to be elevated to an extreme degree due to the adsorption to the container as well as decomposition, thus resulting in a rapid degrading of the quality during storage. Consequently, it is extremely difficult to attain an aqueous preparation stably containing an effective amount of vitamin A. Thus, a vitamin A aqueous preparation which is satisfactory has not been put into practical
45 use.

The inventors made an extensive study to solve the problem. Finally, the inventors have succeeded in solving the thus-far insurmountable problem and produced a stable aqueous preparation which can be used as an eye drop by incorporating, in materializing a vitamin A preparation, a buffering component selected from the group consisting of phosphoric acid, boric acid, citric acid, glutamic acid, ϵ -aminocaproic acid and
50 the alkali metal salts thereof, a non-ionic surfactant and a chelating agent, and adjusting the acidity in the range of pH 5 to pH 8, and filling into a tight container made of polypropylene or polyethylene terephthalate, wherein said tight container is enclosed, with an oxygen scavenger, in another container which is substantially impermeable to oxygen.

SUMMARY OF THE INVENTION

The present invention, therefore, is an aqueous preparation containing a vitamin A derivative which may be used for the treatment of dry eye syndrome characterized in that it is a stable aqueous preparation

containing a buffering component selected from the group consisting of phosphoric acid, boric acid, citric acid, glutamic acid, ϵ -aminocaproic acid and the alkali metal salts thereof, a non-ionic surfactant and a chelating agent, having the acidity adjusted in the range of pH 5 to pH 8, and filled in a tight container made of polypropylene or polyethylene terephthalate, wherein said tight container is enclosed, with an oxygen scavenger, in another container which is substantially impermeable to oxygen.

DETAILED DISCUSSION

The main active constituent of the aqueous preparation of the present invention is vitamin A (retinol) or a vitamin A derivative having vitamin A activities. The vitamin A derivatives having vitamin A activity include, for example, esters of vitamin A such as retinol palmitate and retinol acetate, and any vitamin A derivatives having vitamin A activities may conveniently be used as the active constituent of the aqueous preparation of the present invention.

The concentration of vitamin A in the aqueous preparation of the present invention may be properly settled in accordance with the purpose of use of the preparation. In the case where it is used as an eye drop for the treatment of dry eye syndrome, vitamin A or its derivative having vitamin A activities as mentioned above may usually be incorporated at a concentration in the range of 50 I.U./ml to 20000 I.U./ml, preferably 200 to 3000 I.U./ml, most preferably 500 to 2000 I.U./ml ("I.U." indicates the vitamin A international unit, which is followed hereinafter).

The present inventors carried out an intensive study to materialize an aqueous preparation containing vitamin A and having the stability which may allow the preparation to be practically used as a pharmaceutical preparation.

As a result, it was found that an aqueous preparation which is stable enough in the acidity range adapted to be used as an eye drop, i.e. pH 5 to pH 8, may be obtained by emulsifying vitamin A in water in the presence of a non-ionic surfactant and then adding a buffering agent selected from the group consisting of phosphoric acid, boric acid, citric acid, glutamic acid, ϵ -aminocaproic acid and the alkali metal salts thereof. An additional extensive study made it clear that the stability of the preparation can be further improved by the addition of a chelating agent. However, the aqueous preparation of vitamin A thus obtained was not sufficiently improved concerning the adsorption of vitamin A to the container, which was commonly observed in vitamin A preparations. Thus, a further improvement in compatibility with the container was needed. The present inventors carried out a further study for a container which may not adsorb vitamin A contained in the preparation obtained above, and found out that the adsorption of vitamin A to the container was substantially prevented only when a container made of polypropylene or polyethylene terephthalate was utilized. On the other hand, it was known that vitamin A is readily oxidized by oxygen in the air. Thus, it was found that the vitamin A contained underwent oxidation during storage and that the rapid lowering of its content during storage could not be prevented. Therefore, a study was made in search of a method which may allow a long-term storage of the aqueous preparation. As a result, a preparation which may stand a long-term storage was successfully obtained by enclosing, with an oxygen scavenger, the above preparation, which was filled in a container made of polypropylene or polyethylene terephthalate, in another container substantially impermeable to oxygen.

In the aqueous preparation of the present invention, a non-ionic surfactant is used in order to form an emulsion by dispersing vitamin A in the aqueous medium. For a non-ionic surfactant, any of the non-ionic surfactants which are conventionally used as constituents of eye drops may be conveniently used. For example, either of polysorbate 80 and polyoxyethylenes hydrogenated castor oil may advantageously be used. The ratio of the content of the non-ionic surfactant to that of vitamin A in the preparation may be suitably selected according to the purpose of use and the way of administration, although it necessarily has to be large enough to form a stable emulsion of vitamin A in the aqueous preparation. When used as an eye drop for the treatment of the dry eyes, the non-ionic surfactant is preferably added in the equivalent or more amount to that of vitamin A. Particularly preferably, the non-ionic surfactant may be used in the range of 1.5 to 5 parts by weight, preferably 2 to 4 parts by weight in respect to 1 part by weight of vitamin A.

In the aqueous preparation of the present invention is added a buffering agent selected from the group consisting of boric acid, citric acid, glutamic acid, ϵ -aminocaproic acid and their alkali metal salts. For alkali metal salts, sodium salts and potassium salts may be conveniently used. When necessary, the buffering agents may be used in combination of two or more of them.

A chelating agent may be added in the aqueous preparation of the present invention. For a chelating agent, any of the chelating agents which are conventionally used as constituents of eye drops may be conveniently used. For example, sodium edetate and sodium citrate may be advantageously used. The amount of the chelating agent to be used may usually be in a range of about 0.001 % to about 0.1 % of the

total amount of the aqueous preparation. It has been found that the addition of a chelating agent not only sharply improves the stability of the aqueous preparation of the present invention but also increases the effect of the preservative, as will be mentioned later.

In the aqueous preparation of the present invention, a preservative may be added as in the cases of usual aqueous preparations so as to prevent any growth of microorganisms after opening the container. For preservatives to be added, the preservatives of the type of quaternary ammonium salts such as benzalkonium chloride and of the type of chlorhexidines such as chlorhexidine digluconate may advantageously be used. Other preservatives also may naturally be used insofar as they meet the object of the present invention. It is known that the effect of the preservatives is often reduced in aqueous preparations containing non-ionic surfactants. However, it has been found that if boric acid or its alkali metal salt is used the reduction of the effect of the preservatives may be prevented even when a non-ionic surfactant is contained in relatively a high amount, as mentioned already. Therefore, the use of boric acid or its alkali metal salts enables to reduce the quantity of preservatives (which might cause a local irritation). In this case, the amount of boric acid or borax (sodium borate) is preferably in the range of 0.5 % to 2.5 %, particularly in the range of 1.0 % to 2.0 %.

It is known that vitamin A, the main constituent of the aqueous preparation of the present invention, may be adsorbed to various containers. For example, it is known that when stored in a container made of polyethylene, vitamin A in the preparation being markedly adsorbed, the concentration of vitamin A in the preparation lowers rapidly. The present inventors made a screening for a container that can minimize the adsorption of vitamin A, and found that adsorption to the container comes to be negligible when a polypropylene or a polyethylene terephthalate container is used. Any of such containers adapted to the use as containers for liquid preparation may usually be utilized conveniently.

As vitamin A is quite susceptible to oxidation as mentioned above, it is still difficult to avoid the influence of oxygen in the air even when a polypropylene or polyethylene terephthalate container is used. Then, investigations were made in search of a countermeasure to this. As a result, it was found that the influence of oxygen can be minimized by enclosing, with an oxygen scavenger, the aqueous preparation, filled in a container mentioned above, in another container which is substantially impermeable to oxygen so as to minimize the effect of oxygen. For a such container substantially impermeable to oxygen, a bag-shaped or tubular container made of a film composed of an oxygen-impermeable polymer membrane laminated with an aluminum foil may preferably be used, although any of sealable containers made of a material substantially impermeable to oxygen may conveniently be used. For example, containers made of polyethylene, polypropylene, polyvinylalcohol, polyvinylidene chloride and the like may also be used likewise.

Besides, unless it opposes the object of the present invention, the aqueous preparation of the present invention may also contain other pharmacologically active ingredients than vitamin A such as vitamin E, and it may also contain isotonicizers such as sodium chloride and glycerol and the like when they are required according to the purpose of use. Thus, the inventors have succeeded in the development of an aqueous preparation containing vitamin A as a main ingredient, the preparation thus far considered hardly possible to develop for practical use due to the adsorption to the container and rapid degradation in aqueous preparations.

Effect of the invention:

The test results concerning the aqueous preparation containing vitamin A as a main constituent, which is provided by the present invention, will be described below to show the effect of the present invention.

1. pH and stability of the aqueous preparation

Vitamin A palmitate was emulsified with a borate buffer system and stored for 1 week at 50 °C and 40 °C to examine the change in vitamin A content in the aqueous preparation. The following results were obtained.

When stored at 50 °C, the remaining rate in the samples of pH 4.0 and pH 8.0 was 83.9 % and 86.4 %, respectively, whereas the remaining rate was more than 90 % in any of the samples which had a pH value between 5.0 and 7.5. On the other hand, when stored at 40 °C, the remaining rate in the samples having a pH value between 5.0 and 7.0 was 95 to 96 %, indicating a notable stability in comparison with the remaining rate in the samples of pH 4.0 and pH 9.0, i.e. about 90 %. The appearance of the emulsion of these samples remained unchanged.

2. Effect of buffer system on stability

For evaluation of buffer systems, the aqueous preparations of pH 5.8 were prepared using different buffer systems based on sodium acetate, borax, citric acid, ϵ -aminocaproic acid, sodium glutamate, sodium hydrogen phosphate and taurine. They were stored for 2 weeks at 40 °C under protection from light with aluminum foil, and measured for the remaining rate. The following results were obtained.

The remaining rate was 81 to 84 % in sodium acetate buffer system and taurine buffer system, and about 88 % in phosphate buffer system. In any other buffer systems, i.e. borax, citric acid, ϵ -aminocaproic acid, and sodium glutamate, the remaining rate was at least 90 %. Thus, the latter, and the phosphate buffer system, were shown to have sufficient stability for use. No change in appearance was observed in any of the samples.

3. Effect of non-ionic surfactant

Polysorbate 80 was applied as a non-ionic surfactant, and the samples were prepared with and without it to compare their stability. However, it was found that any emulsified state could not be attained in the sample without polysorbate 80. In addition, it was also found that the degradation of vitamin A in the aqueous preparation became rapid when a non-ionic surfactant with a high peroxide value was used.

4. Effect of chelating agent

Sodium edetate was applied as a chelating agent, and aqueous preparations containing it at concentrations of 0.005 %, 0.01 %, 0.05 % and 0.1 % were prepared. These were stored for 6 weeks at 40 °C and 30 °C with an aqueous preparation containing no sodium edetate. The following results were obtained.

Any of above samples exhibited a sufficient stability, i.e. the remaining rate of vitamin A was not less than 93 % at 40 °C, and not less than 97 % at 30 °C. In the sample without sodium edetate, the rate was 90 % at 40 °C and 96 % at 30 °C. The result thus indicated that it is preferable to add sodium edetate in an amount of not less than 0.005 % in order to obtain a stable aqueous preparation.

5. Study on container materials

Each cut piece of containers for eye drops made of polyethylene, polyethylene terephthalate or polypropylene was put in each glass ampoule, and the aqueous preparation of the present invention was added to it. After stored for 4 days at 60 °C or for 14 days at 50 °C, the remaining rate of vitamin A was determined. The results were as follows.

While the remaining rate of vitamin A was lowered to 77 % after the 4-day storage at 60 °C due to the adsorption of vitamin A to the container material in the case of polyethylene container, the remaining rate of not less than 80 % was observed in other samples. In the test at 50 °C for 14 days, the remaining rate was lowered nearly to 60 % in the case of polyethylene, whereas the remaining rate of not less than 70 % was observed in other samples, indicating that the latter can be aqueous preparations which allow practical use.

6. Study on oxygen scavenger

The aqueous preparation of the present invention was filled in a polypropylene container, and then put in a bag made of laminated aluminum film together with an oxygen scavenger [from MITSUBISHI GAS KAGAKU; AGELESS (trademark) Z-30] to test the stability. While the remaining rate of vitamin A was lowered to about 80 % after 6-month storage at room temperature in the case of the sample without the oxygen scavenger, the remaining rate of not less than 95 % was observed in the case where the oxygen scavenger was used. In the sample where an oxygen scavenger was used, 93 % of vitamin A was found remained after 1 week even under a severe condition of 50 °C.

7. Effect of the preservative

The test of the preservative effect was carried out according to the method of preservative test in the U.S. Pharmacopeia XXI. The following strains of microorganisms were selected for the test.

Staphylococcus aureus	IFO	13276
Escherichia coli	IFO	3972
Pseudomonas aeruginosa	IFO	13275
Candida albicans	IFO	1594
Aspergillus niger	IFO	9455

The formula of the aqueous preparation used in this test was as follows.

retinol palmitate	200000 I.U.
boric acid	1.7 g
sodium edetate	0.01 g
polysorbate 80	0.85 g
benzalkonium chloride (10 %) solution	0.05 ml
borax	q.s.
<u>sterile purified water</u>	<u>to 100 ml</u>
	pH 7.0

The result of the test showed that none of these microorganisms survived on the 14th day after inoculation. Staphylococcus, Escherichia and Pseudomonas, in particular, were not found surviving on the 7th day after inoculation. Thus, it was demonstrated that the aqueous preparation of the present invention has a sufficient preservative effect for the use as an eye drop.

Examples

Example 1

5

retinol palmitate 50000 I.U.

polysorbate 80 0.17 g

10

borax 0.01 g

boric acid 1.7 g

15

sodium edetate 0.01 g

benzalkonium chloride
(10 %) solution 0.05 ml

20

sterile purified water to 100 ml

pH 5.8

Example 2

25

retinol palmitate 150000 I.U.

polysorbate 80 0.51 g

30

borax 0.01 g

boric acid 1.7 g

sodium edetate 0.01 g

35

benzalkonium chloride
(10 %) solution 0.05 mlsterile purified water to 100 ml

40

pH 7.0

45

50

55

Example 3

5	retinol palmitate	200000	I.U.
	polysorbate 80	0.85	g
	borax	0.01	g
10	boric acid	1.7	g
	sodium dihydrogen phosphate	0.2	g
15	sodium chloride	0.9	g
	sodium edetate	0.005g	
20	benzalkonium chloride (10 %) solution	0.05	ml
	hydrochloric acid	q.s.	
	<u>sterile purified water</u>	<u>to 100</u>	<u>ml</u>
25		pH 7.0	

Example 4

30	retinol palmitate	200000	I.U.
	polyoxyethylenehydrogenated castor oil 60	0.5	g
35	glycerol	2.6	g
	citric acid	0.2	g
40	chlorhexidine gluconate (20%) solution	0.025	ml
	sodium edetate	0.005	g
	sodium hydroxide	q.s.	
45	<u>sterile purified water</u>	<u>to 100</u>	<u>ml</u>
		pH 7.0	

50

55

Example 5

5	retinol palmitate	100000	I.U.
	polyoxyethylenehydrogenated castor oil 60	0.2	g
10	glycerol	0.5	g
	borax	0.2	g
	boric acid	1.5	g
15	benzalkonium chloride (10 %) solution	0.05	ml
	sodium edetate	0.01	g
20	sterile purified water	to 100	ml
		pH 7.0	

Example 6

25	retinol palmitate	50000	I.U.
	polyoxyethylenehydrogenated castor oil 60	0.1	g
30	glycerol	0.5	g
	borax	0.2	g
35	boric acid	1.5	g
	benzalkonium chloride (10 %) solution	0.05	ml
40	sodium edetate	0.01	g
	sterile purified water	to 100	ml
45		pH 7.0	

Claims

50 Claims for the following Contracting States : AT, BE, CH, DE, DK, FR, GB, IT, LI, LU, NL, SE

1. A stable aqueous preparation for ophthalmic topical administration containing vitamin A or a vitamin A derivative having vitamin A activities to be used for the treatment of dry eye syndrome characterized in that it contains a buffering component selected from the group consisting of phosphoric acid, boric acid, citric acid, glutamic acid, ϵ -aminocaproic acid and the alkali metal salts thereof, a non-ionic surfactant and a chelating agent, that its acidity is adjusted in the range of pH 5 to pH 8, and that it is filled in a tight container made of polypropylen or polyethylene terephthalate, wherein said tight container is enclosed, with an oxygen scavenger, in another container which is substantially imperme-

able to oxygen.

2. The stable aqueous preparation of Claim 1 wherein the vitamin A derivative is retinol palmitate.
- 5 3. The stable aqueous preparation of Claim 1 or 2, wherein the non-ionic surfactant is polysorbate 80 or polyoxyethylenehydrogenated castor oil 60.
4. The stable aqueous preparation of Claim 1, 2 or 3, wherein the chelating agent is sodium edetate.
- 10 5. The stable aqueous preparation of Claim 1, 2, 3 or 4, which contains a quaternary ammonium salt or a salt of chlorhexidine as a preservative.
6. The stable aqueous preparation of Claim 5, wherein the quaternary ammonium salt is benzalkonium chloride.
- 15 7. The stable aqueous preparation of Claim 5, wherein the salt of chlorhexidine is chlorhexidine digluconate.

Claims for the following Contracting States : ES, GR

- 20 1. A method for preparing a stable aqueous preparation for ophthalmic topical administration containing vitamin A or a vitamin A derivative having vitamin A activities to be used for the treatment of dry eye syndrome, characterized by providing a buffering component selected from the group consisting of phosphoric acid, boric acid, citric acid, glutamic acid, ϵ -aminocaproic acid and the alkali metal salts thereof, a non-ionic surfactant and a chelating agent; and adjusting the acidity in the range of pH 5 to pH 8; wherein the components are filled in a tight container made of polypropylene or polyethylene terephthalate, and said tight container is enclosed, with an oxygen scavenger, in another container which is substantially impermeable to oxygen.
- 25 2. The method of Claim 1 wherein the vitamin A derivative is retinol palmitate.
3. The method of Claim 1 or 2, wherein the non-ionic surfactant is polysorbate 80 or polyoxyethylenehydrogenated castor oil 60.
- 35 4. The method of Claim 1, 2 or 3, wherein the chelating agent is sodium edetate.
5. The method of Claim 1, 2, 3 or 4, wherein the preparation contains a quaternary ammonium salt or a salt of chlorhexidine as a preservative.
- 40 6. The method of Claim 5, wherein the quaternary ammonium salt is benzalkonium chloride.
7. The method of Claim 5, wherein the salt of chlorhexidine is chlorhexidine digluconate.

45 Patentansprüche

Patentansprüche für folgende Vertragsstaaten : AT, BE, CH, DE, DK, FR, GB, IT, LI, LU, NL, SE

- 50 1. Stabile, wäßrige Zubereitung zur ophthalmischen, topischen Verabreichung, enthaltend Vitamin A oder ein Vitamin A-Derivat mit Vitamin A-Aktivität, zur Verwendung bei der Behandlung des Trockene-Augen-Syndroms, dadurch gekennzeichnet, daß sie eine Pufferkomponente, ausgewählt unter Phosphorsäure, Borsäure, Zitronensäure, Glutaminsäure, ϵ -Aminocaprinsäure und den Alkalimetallsalzen davon, ein nicht-ionisches, oberflächenaktives Mittel und ein chelatbildendes Mittel enthält, daß deren Acidität eingestellt ist auf einem Wert im Bereich von pH 5 bis pH 8, und daß sie in einem dichten Behälter, hergestellt aus Polypropylen oder Polyethylen-Terephthalat, abgefüllt ist, wobei der dichte Behälter zusammen mit einem Sauerstofffänger in einem anderen Behälter eingeschlossen ist, der im wesentlichen impermeabel ist für Sauerstoff.
- 55 2. Stabile, wäßrige Zubereitung nach Anspruch 1, wobei das Vitamin A-Derivat Retinolpalmitat ist.

3. Stabile, wäßrige Zubereitung nach Anspruch 1 oder 2, worin das nicht-ionische, oberflächenaktive Mittel Polysorbat 80 oder Polyoxyethylen-hydrogeniertes Castoröl 60 ist.
- 5 4. Stabile, wäßrige Zubereitung nach Anspruch 1, 2 oder 3, worin das chelatbildende Mittel Natriumedetat ist.
5. Stabile, wäßrige Zubereitung nach Anspruch 1, 2, 3 oder 4, die ein quaternäres Ammoniumsalz oder ein Chlorhexidin-Salz als Konservierungsmittel enthält.
- 10 6. Stabile, wäßrige Zubereitung nach Anspruch 5, worin das quaternäre Ammoniumsalz Benzalkoniumchlorid ist.
7. Stabile, wäßrige Zubereitung nach Anspruch 5, worin das Chlorhexidinsalz Chlorhexidin-Digluconat ist.

15 **Patentansprüche für folgende Vertragsstaaten : ES, GR**

1. Verfahren zur Herstellung einer stabilen, wäßrigen Zubereitung für die ophthalmische topische Verabreichung, enthaltend Vitamin A oder ein Vitamin A-Derivat mit Vitamin A-Aktivität, zur Verwendung bei der Behandlung des Trockene-Augen-Syndroms,
20 dadurch gekennzeichnet, daß man eine Pufferkomponente, ausgewählt unter Phosphorsäure, Borsäure, Zitronensäure, Glutaminsäure, ϵ -Aminocaprinsäure und den Alkalimetallsalzen davon, ein nicht-ionisches, oberflächenaktives Mittel und ein chelatbildendes Mittel bereitstellt; und deren Acidität auf einen Wert im Bereich von pH 5 bis pH 8 einstellt; wobei die Komponenten in einem dichten Behälter, hergestellt aus Polypropylen oder Polyethylen-Terephthalat, abgefüllt werden, wobei der dichte Behälter
25 zusammen mit einem Sauerstofffänger in einem anderen Behälter eingeschlossen wird, der im wesentlichen impermeabel ist für Sauerstoff.
2. Verfahren nach Anspruch 1, wobei das Vitamin A-Derivat Retinolpalmitat ist.
- 30 3. Verfahren nach Anspruch 1 oder 2, worin das nichtionische, oberflächenaktive Mittel Polysorbat 80 oder Polyoxyethylen-hydrogeniertes Castoröl 60 ist.
4. Verfahren nach Anspruch 1, 2 oder 3, worin das chelatbildende Mittel Natriumedetat ist.
- 35 5. Verfahren nach Anspruch 1, 2, 3 oder 4, wobei das Präparat ein quaternäres Ammoniumsalz oder ein Chlorhexidin-Salz als Konservierungsmittel enthält.
6. Verfahren nach Anspruch 5, worin das quaternäre Ammoniumsalz Benzalkoniumchlorid ist.
- 40 7. Verfahren nach Anspruch 5, worin das Chlorhexidinsalz Chlorhexidin-Digluconat ist.

Revendications

Revendications pour les Etats contractants suivants : AT, BE, CH, DE, DK, FR, GB, IT, LI, LU, NL, SE

- 45 1. Préparation aqueuse stable pour l'administration locale ophtalmique contenant de la vitamine A ou un dérivé de la vitamine A ayant des activités de vitamine A, destinée à l'utilisation pour le traitement du syndrome de l'oeil sec, caractérisée en ce qu'elle contient un constituant tampon choisi parmi l'acide phosphorique, l'acide borique, l'acide citrique, l'acide glutamique, l'acide ϵ -aminocaproïque et leurs sels de métaux alcalins, un agent tensioactif non-ionique et un agent chélatant, en ce que son acidité est
50 ajustée dans l'intervalle de pH 5 à pH 8 et en ce qu'elle est introduite dans un récipient hermétique en polypropylène ou en polyéthylène téréphtalate, ledit récipient hermétique étant enfermé, avec un absorbeur d'oxygène, dans un autre récipient qui est pratiquement imperméable à l'oxygène.
2. Préparation aqueuse stable selon la revendication 1, caractérisée en ce que le dérivé de la vitamine A est le palmitate de rétinol.
- 55 3. Préparation aqueuse stable selon les revendications 1 ou 2, dans laquelle l'agent tensio-actif non-ionique est le polysorbate 80 ou l'huile de ricin hydrogénée-polyoxyéthylène 60.

4. Préparation aqueuse stable selon les revendications 1, 2 ou 3, dans laquelle l'agent chélatant est l'édétate de sodium.
5. Préparation aqueuse stable selon les revendications 1, 2, 3 ou 4, qui contient un sel d'ammonium quaternaire ou un sel de chlorhexidine comme conservateur.
6. Préparation aqueuse stable selon la revendication 5, dans laquelle le sel d'ammonium quaternaire est le chlorure de benzalkonium.
7. Préparation aqueuse stable selon la revendication 5, dans laquelle le sel de chlorhexidine est le digluconate de chlorhexidine.

Revendications pour les Etats contractants suivants : ES, GR

1. Procédé d'obtention d'une préparation aqueuse stable pour l'administration locale ophtalmique contenant de la vitamine A ou un dérivé de la vitamine A ayant des activités de vitamine A, destinée à l'utilisation pour le traitement du syndrome de l'oeil sec, caractérisé en ce qu'on apporte un constituant tampon choisi parmi l'acide phosphorique, l'acide borique, l'acide citrique, l'acide glutamique, l'acide ϵ -aminocaproïque et leurs sels de métaux alcalins, un agent tensio-actif non-ionique et un agent chélatant; et en ce qu'on ajuste l'acidité dans l'intervalle de pH 5 à pH 8; dans lequel les constituants sont introduits dans un récipient hermétique en polypropylène ou en polyéthylène téréphtalate, et ledit récipient hermétique étant enfermé, avec un absorbeur d'oxygène, dans un autre récipient qui est pratiquement imperméable à l'oxygène.
2. Procédé selon la revendication 1, dans lequel le dérivé de la vitamine A est le palmitate de rétinol.
3. Procédé selon les revendications 1 ou 2, dans lequel l'agent tensio-actif non-ionique est le polysorbate 80 ou l'huile de ricin hydrogénée-polyoxyéthylène 60.
4. Procédé selon les revendications 1, 2 ou 3, dans lequel l'agent chélatant est l'édétate de sodium.
5. Procédé selon les revendications 1, 2, 3 ou 4, dans lequel la préparation contient un sel d'ammonium quaternaire ou un sel de chlorhexidine comme conservateur.
6. Procédé selon la revendication 5, dans lequel le sel d'ammonium quaternaire est le chlorure de benzalkonium.
7. Procédé selon la revendication 5, dans lequel le sel de chlorhexidine est le digluconate de chlorhexidine.